

L1063503 ZAMB/06Y PPE 4122625



Fusilade Forte[®]

150EC

syngenta.

**TO CAUSE A HAZARD IN THE USE, STORAGE
OR DISPOSAL OF THIS SUBSTANCE IS AN OFFENCE.**

Before opening the container, read the safety advice.

Hazard Classification:

WHO Class II (moderately hazardous).

Shelf-Life: At least 2 years from date of formulation,
in its original unopened container.

Emergency Call Number:

Swiss Toxicological Information Centre,
(24 h) +41 44 251 51 51

Manufacturer:

Syngenta Crop Protection AG, Basel, Switzerland

Registered and distributed by:

Syngenta Zambia Ltd.

Plot 5255, Mukwa Road,

P.O. Box 33088,

Lusaka, Zambia,

Tel.: +260 211 244 766



5 Litres

Product names marked ® or TM, the ALLIANCE FRAME
the SYNGENTA Logo and the PURPOSE ICON
are Trademarks of a Syngenta Group Company



WARNING



DIRECTIONS FOR USE

1. Fusilade Forte® 150 EC AS A HERBICIDE:

Fusilade Forte® 150 EC may be used to control most annual and perennial grass weeds as well as the suppression of Upright Starbur in Burley and Virginia tobacco in the lands, cotton, groundnuts, beans and soyabeans.

PRECAUTIONS:

- Handle with care; avoid splashing / contact; poisonous by swallowing, inhalation and skin contact. Do not smoke, eat or drink while using this product.
- Store in original container under LOCK AND KEY.
KEEP OUT OF REACH OF CHILDREN.
KEEP APART FROM FOOD AND FOODSTUFFS.
- MAY BE DANGEROUS TO ANIMALS, BIRDS, FISH AND BEES.
- DO NOT CONTAMINATE DRINKING POOLS, DAMS, RIVERS, WATERWAYS AND OTHER WATER SUPPLIES.

FOR SAFETY WHEN MIXING

- Wear eye protection (at least glasses). Undiluted Fusilade Forte® 150 EC can irritate your eyes. If you get it in your eyes, wash it out at once. **See FIRST AID.**
- Wear rubber gloves. Undiluted Fusilade Forte® 150 EC can irritate your skin. If you get it on your skin, wash it off at once. If you spill it on your clothes, change and wash them immediately.
- Triple rinse empty container and pour washings into spray tank.

FOR SAFETY DURING SPRAYING:

- As with all chemicals AVOID CONTACT WITH SPRAY as much as you can.
- Wear overalls and boots.

FOR SAFETY AFTER SPRAYING:

- Dispose of unwanted spray solution by spraying off on waste ground.
- Remove protective clothing and wash it before re-use.
- WASH YOURSELF with soap and water.
- Destroy empty container – DO NOT USE IT for any other purpose. Crush or cut it, then burn or bury it.
- Clean applicator thoroughly.
- Dispose of wash water in a safe manner.

SYMPTOMS OF HUMAN POISONING:

No known symptoms of poisoning.

FIRST AID:

Eye splashes: Hold eyelids apart and pour in a gentle stream of water for 10-15 minutes. Go to a doctor. In case of contact with skin, remove contaminated clothing; wash affected body parts immediately with plenty of water and soap.

If Fusilade Forte® 150 EC is SWALLOWED do not make the person vomit.

Take person, and this label, to a DOCTOR AT ONCE.

NOTE TO PHYSICIAN:

Perform gastric lavage, taking care to prevent aspiration of gastric contents. Consider administration of activated charcoal and a laxative.

Treat symptomatically.

RATES

| CROP STAGE | WEEDS/WEED HEIGHT | RATE L / HA |
|---|--|-------------|
| Tobacco Seedbed | Annual grasses | 0.58 |
| Field (tobacco, cotton, groundnuts, beans and soyabeans) | Annual grasses: < 120 mm | 0.830 |
| | 120-250 mm | 1.25 |
| | > 250 mm | 1.7-2.1 |
| | Mature perennial grasses: Undisturbed rhizomes | 2.5-6.7 |
| | Fragmented rhizomes | 1.25-2.9 |
| | Shamva and sorghum volunteers (up to 200 mm height) | 0.4 |
| | Wheat and barley volunteers (up to 5 leaf stage) | 1.25 |

APPLICATION:

A relatively short contact period of 1.5-2 hours without rain is required for satisfactory results.

Tobacco: Fusilade Forte® 150 EC can be directed onto grass weeds or applied as a broadcast spray anytime after transplanting, until within 3 weeks of harvesting.

Fusilade Forte® 150 EC contains a wetter.

No additional wetter is required.

Fusilade Forte® 150 EC may be applied through any sprayer fitted with normal fan, cone or flood jet nozzles.

For ground application 100-400 litres spray / ha should be applied.

For aerial application 20-30 litres spray / ha should be applied.

Clean water should always be used.

HARVEST INTERVAL

(Providing rate has not exceeded 3.0 litres/ha):

Cotton, Groundnuts, Phaseolus beans and Soyabeans: 8 weeks

Tobacco: 3 weeks

WARNINGS:

Do not plant any of the following crops: maize, sorghum, wheat, barley, or grasses within 2 months of a Fusilade Forte® 150 EC application.

With Fusilade Forte® 150 EC, weeds should be actively growing and not stressed, at the time of application.

COMPATIBILITY:

It is not recommended to tank mix Fusilade Forte® 150 EC with other materials.

These should be sprayed in a sequence of at least five days apart. The order in which Fusilade Forte® 150 EC and other products are applied is not important.

2. Fusilade Forte® 150 EC AS A SUGAR CANE RIPENER: APPLICATION DIRECTIONS AND RATES:

Fusilade Forte® 150 EC should be applied to EARLY season sugar cane harvests (April-June harvests).

| APPLICATION | RATE | COMMENT |
|--------------|------------|--|
| Early season | 275 ml /ha | The time of application is related to the date of harvest (i.e. 6-10 weeks prior to the set harvesting date) however, application also depends on cane maturity. |

WARNING:

Spray drift or direct contact with young cane can result in crop damage.

SPILLAGE AND LEAKAGE

Remove heavily contaminated soil layer and bury in a safe place away from water supplies. Collect up spilt material thoroughly and bury it in a safe place or put it in a waste bin or in a plastic bag and dispose of on a landfill site approved for pesticides.

Rinse spill area with plenty of water.

Note: Spilt product must not be re used.

PRODUCT AND CONTAINER DISPOSAL

Provided there are no local legal or other rules and regulations, the following procedure is recommended: Dispose of surplus product on a landfill site approved for pesticides or bury in a safe place away from water supplies. Empty the product sachets / containers well.

Dispose of containers made of cardboard, paper or plastic on a landfill site or burn in a safe place. If burned stay out of the smoke. Be sure that the smoke or fumes will not damage nearby plants, crops or trees, or harm animals.


DO NOT re use empty containers for any other purpose.

User's risk

The user bears the risk for damage resulting from factors beyond the manufacturer's control.

Resistant strains of weeds may develop or may exist against which herbicides may be less effective. Since the occurrence of such strains cannot be forecast, neither the manufacturer nor its distributors can assume responsibility for any loss or damage caused by failure to control resistant weeds.

All recommendations for use of the herbicide are based on the current state of the manufacturer's knowledge. Since the manufacturer cannot control the farmers' choice of herbicides, its application, use, storage or processing of the product or other agronomic practices, the manufacturer cannot assume responsibility therefor.



NO TEXT HERE
GLUE PAGE
97 x 134 mm